



---

**Anterior Cruciate Ligament Reconstruction: Accelerated Rehabilitation Protocol\***

---

This rehabilitation protocol was developed for patients who have anterior cruciate ligament (ACL) reconstruction and desire to return to strenuous sports activities early postoperatively.

The following are **exclusionary criteria** for this protocol:

- v Concomitant meniscal repair
- v Concomitant ligament reconstruction
- v Concomitant patellofemoral realignment procedure
- v ACL revision reconstruction
- v MRI evidence of bone bruising or articular cartilage damage seen during procedure

The protocol is divided into 7 phases according to postoperative weeks (for instance, Phase I = Postoperative Weeks 1-2). Each phase has several categories including:

- v *General observation* of the patient's condition (weight bearing, pain, hemarthrosis, muscle control)
- v *Evaluation* of specific variables with *goals* identified for each
- v Treatment and exercise program, according to *frequency* and *duration*
- v *Rehabilitation goals* which must be achieved to enter into the next phase

The **overall goals** of the reconstruction and rehabilitation are to:

- v Control joint pain, swelling, hemarthrosis (minimal or none)
- v Regain normal knee flexion and extension
- v Regain a normal gait pattern and neuromuscular stability for ambulation
- v Regain normal quadriceps, hamstring lower extremity muscle strength
- v Regain normal proprioception, balance, and coordination for desired activities
- v Achieve optimal functional outcome based on orthopaedic and patient goals

The supervised rehabilitation program is supplemented with a *home self-management program* which the patient performs on a daily basis. The therapist must evaluate the patient thoroughly to implement the enclosed protocol and should see the patient in the clinic for therapeutic procedures and modality treatments which are required for rehabilitation. The majority of this protocol can be accomplished at home provided patient cooperation and follow through are present. The approximate number of rehabilitation visits required for each phase are provided. Additional supervision may be required if a complication develops.

**Important postoperative signs** to monitor include:

- v Swelling of the knee joint or soft tissues (avoid hemarthrosis)
- v Abnormal pain response
- v Abnormal gait pattern with or without assistive device
- v Insufficient flexion or extension motions, limited patellar mobility
- v Weakness (strength/control) of the lower extremity, especially the quads/hamstrings
- v Insufficient lower extremity flexibility
- v Abnormal AP displacement (joint arthrometer testing > 3 mm) early postoperatively

The patient is placed into one of four **sports activity** and **occupational activity categories** based on the following scales. It is expected that patients who follow this protocol desire to return to sports activity levels I or II, or very heavy/heavy occupations.

### 1. Cincinnati Knee Rating System Sports Activity Scale

(check one)

0 Level I - jumping, hard pivoting, cutting sports (basketball, volleyball, football, gymnastics, soccer)

0 Level II - running, twisting, turning (tennis, racquetball, handball, ice/field hockey, skiing, wrestling)

0 Level III - light recreational sports (bicycling, swimming - no running, twisting, jumping)

0 Level IV - no sports, activities of daily living only

### 2. Cincinnati Knee Rating System Occupational Rating Scale

Factor 1 sitting	Factor 2 standing/ walking	Factor 3 walking on uneven ground	Factor 4 squatting	Factor 5 climbing	Factor 6 lifting/ carrying	Factor 7 pounds carried
0 0 8-10 hrs/day	0 0 0 hrs/day	0 0 0 hrs/day	0 0 0 times/day	0 0 0 times/day	0 0 0 times/day	0 0 0-5 lbs
1 0 6-7 hrs/day	2 0 1 hrs/day	2 0 1 hrs/day	1 0 1-5 times/day	2 0 1 flight 2 times/day	1 0 1-5 times/day	1 0 6-10 lbs
2 0 4-5 hrs/day	4 0 2-3 hrs/day	4 0 2-3 hrs/day	2 0 6-10 times/day	4 0 3 flights 2 times/day	2 0 6-10 times/day	2 0 11-20 lbs
3 0 2-3 hrs/day	6 0 4-5 hrs/day	6 0 4-5 hrs/day	3 0 11-15 times/day	6 0 10 flights/ ladders	3 0 11-15 times/day	3 0 21-25 lbs
4 0 1 hrs/day	8 0 6-7 hrs/day	8 0 6-7 hrs/day	4 0 16-20 times/day	8 0 ladders with weight 2-3 days/week	4 0 16-20 times/day	4 0 26-30 lbs
5 0 0 hrs/day	10 0 8-10 hrs/day	10 0 8-10 hrs/day	5 0 > 20 times/day	8 0 ladders daily with weight	5 0 > 20 times/day	5 0 > 20 lbs

\_\_\_\_\_ points x 2 = \_\_\_\_\_ total points

#### Occupation Rating

#### Total Points

0 Disabled	0
0 Very light	1-20
0 Light	21-40
0 Moderate	41-60
0 Heavy	61-80
0 Very heavy	> 80

## Physical Therapy Visit Timeline\*

Phase	Weeks Postoperative	Minimum # Visits	Maximum # Visits
1	1-2	2	4
2	3-4	2	4
3	5-6	1	2
4	7-8	1	2
5	9-12	1	2
6	13-26	2	3
7	27-52	2	4
Total		11	21

### \*Physician Notification

The physician will be notified if the patient (1) fails to meet the expected goals for each phase of the protocol, (2) has a persistent joint effusion, (3) develops a chronic pain syndrome, (4) develops abnormal AP displacement on arthrometer testing (> 3 mm difference), (5) complains of giving-way, (6) has difficulty with ambulation, or (7) has a limitation of knee motion or patellar mobility.

These problems could result in a modification of this protocol and necessitate further visits to the physical therapist.

### Discharge Criteria (If goals for sports and occupation place patient into different categories, use category with highest functional demand criteria. For symptoms, use Symptom Rating Form found on next page.)

#### *Sports Activity Level I or Heavy/Very Heavy Occupational Rating\**

No pain, swelling, giving-way with level 10 on Symptom Rating Form

KT-2000 < 3 mm (I-N, 134 N, total A/P)

Biodex strength testing: quadriceps & hamstrings strength  $\geq$  85% of opposite limb

Function testing: 2 hop tests, limb symmetry  $\geq$  85%

#### *Sports Activity Level II or Moderate Occupational Rating\**

No pain, swelling, giving-way with level 8 on Symptom Rating Form

KT-2000 < 3 mm (I-N, 134 N, total A/P)

Biodex strength testing: quadriceps & hamstrings strength  $\geq$  80% of opposite limb

Function testing: 2 hop tests, limb symmetry  $\geq$  85%

#### *Sports Activity Level III or Light Occupational Rating*

No pain, swelling, giving-way with level 6 on Symptom Rating Form

KT-2000 3-5 mm (I-N, 134 N, total A/P)

Biodex strength testing: quadriceps & hamstrings strength  $\geq$  70% of opposite limb

Function testing: 2 hop tests, limb symmetry  $\geq$  75%

#### *Sports Activity Level IV (ADL) or Very light Occupational Rating*

No pain, swelling, giving-way with level 4 on Symptom Rating Form

KT-2000 3-5 mm (I-N, 134 N, total A/P)

Biodex strength testing: quadriceps & hamstrings strength < 70% of opposite limb

Function testing: 2 hop tests, limb symmetry < 75%

\*patients desiring to return to sports or strenuous work activities may require 4-6 more physical therapy visits during postoperative weeks 25-52 for advanced neuromuscular, strength, and activity-specific training to prevent reinjury.

# Cincinnati Knee Rating System Symptom Rating Form

Scale	Description
10	Normal knee, able to do strenuous work/sports with jumping, hard pivoting
8	Able to do moderate work/sports with running, turning and twisting; symptoms with strenuous work/sports
6	Able to do light work/sports with no running, twisting or jumping; symptoms with moderate work/sports
4	Able to do activities of daily living alone; symptoms with light work/sports
2	Moderate symptoms (frequent, limiting) with activities of daily living
0	Severe symptoms (constant, not relieved) with activities of daily living

## 1. PAIN (circle one)

10 — 8 — 6 — 4 — 2 — 0

## 2. SWELLING (circle one)

10 — 8 — 6 — 4 — 2 — 0

## 3. PARTIAL GIVING-WAY (circle one) (partial knee collapse, no fall to the ground)

10 — 8 — 6 — 4 — 2 — 0

## 4. FULL GIVING-WAY (circle one) (knee collapse occurs with actual falling to the ground)

10 — 8 — 6 — 4 — 2 — 0

## Return to Activities Warning

Return to strenuous activities after major knee ligament surgery carries the definite risk of a repeat injury or the potential of compounding the original injury. These risks cannot always be scientifically assessed. Patients are warned to return to athletic activities carefully and to avoid any activity in which pain, swelling, or a feeling of instability is present.

## References

- Heckmann TP, Noyes FR, and Barber-Westin SD: Autogenic and allogeneic anterior cruciate ligament rehabilitation. In *Knee Ligament Rehabilitation*, Ellenbecker TS (ed), Churchill Livingstone, New York, NY, pp. 132-150, 2000
- Noyes FR, Berrios-Torres S, Barber-Westin SD, and Heckmann TP: Prevention of permanent arthrofibrosis after anterior cruciate ligament reconstruction alone or combined with associated procedures: A prospective study in 443 knees. *Knee Surgery, Sports Traumatology, Arthroscopy* 8: 196-206, 2000.
- Barber-Westin SD, Noyes FR, and McCloskey JW: Rigorous statistical reliability, validity, and responsiveness testing of the Cincinnati Knee Rating System in 350 subjects with uninjured, injured, or anterior cruciate ligament-reconstructed knees. *Am J Sports Med.* 27: 402-416, 1999.
- Barber-Westin SD, Noyes FR, Heckmann TP, and Shaffer BL: The effect of exercise and rehabilitation on anterior-posterior knee displacements following anterior cruciate ligament autograft reconstruction. *Am. J. Sports Med.* 27: 84-93, 1999.
- Noyes FR and Barber-Westin SD: A comparison of results in acute and chronic anterior cruciate ligament ruptures of arthroscopic-assisted autogenous patellar tendon reconstruction. *Am J Sports Med* 25: 460-471, 1997.
- Barber-Westin SD, Noyes FR, and Andrews MA: A rigorous comparison between the sexes of results and complications after anterior cruciate ligament reconstruction. *Am. J. Sports Med.* 25: 514-526, 1997.
- Noyes FR, Mooar LA, and Barber SD: The assessment of work-related activities and limitations in knee disorders. *Am J Sports Med* 19: 178-188, 1991.
- Noyes FR, Barber SD, and Mangine RE: Abnormal lower limb symmetry determined by function hop tests after anterior cruciate ligament rupture. *Am J Sports Med* 19: 513-518, 1991.
- Noyes FR, Barber SD, and Mooar LA: A rationale for assessing sports activity levels and limitations in knee disorders. *Clin Orthop Rel Res* 246: 238-249, 1989.



## Cincinnati Sportsmedicine and Orthopaedic Center Rehabilitation Protocol Summary for Accelerated ACL Reconstruction

	Postoperative Weeks					Postoperative Months			
	1-2	3-4	5-6	7-8	9-12	4	5	6	7-12
<b>Brace:</b> immobilizer for patient comfort	X	(X)							
<b>Range of motion minimum goals:</b> 0-110° 0-120° 0-135°	X	X	X						
<b>Weight bearing:</b> 1/2 body weight Full	X	X							
<b>Patella mobilization</b>	X	X	X						
<b>Modalities:</b> Electrical muscle stimulation (EMS) Pain/edema management (cryotherapy)	X X	X X	X X	X X	X X	X X	X X	X X	X X
<b>Stretching:</b> Hamstring, gastroc-soleus, iliotibial band, quadriceps	X	X	X	X	X	X	X	X	X
<b>Strengthening:</b> Quad isometrics, straight leg raises, active knee extension Closed-chain: gait retraining, toe raises, wall sits, mini-squats Knee flexion hamstring curls (90°) Knee extension quads (90-30°) Hip abduction-adduction, multi-hip Leg press (70-10°)	X X X X X X	X X X X X X	X X X X X X	X X X X X X	X X X X X X	X X X X X X	X X X X X X	X X X X X X	X X X X X X
<b>Balance/proprioceptive training:</b> Weight-shifting, cup walking, BBS BBS, BAPS, perturbation training, balance board, mini-trampoline	X	X	X	X X	X X	X X	X X	X X	X X
<b>Conditioning:</b> UBC Bike (stationary) Aquatic program Swimming (kicking) Walking Stair climbing machine Ski machine Elliptical machine	X	X X X	X X X	X X X X	X X X X	X X X X	X X X X	X X X X	X X X X
Running: straight					X	X	X	X	X
Cutting: lateral carioca, figure 8's						X	X	X	X
Plyometric training						X	X	X	X
Full sports							X	X	X

BAPS = Biomechanical Ankle Platform System (Camp, Jackson, MI), BBS = Biodex Balance System (Biodex Medical Systems, Inc, Shirley, NY), UBC = upper body cycle (Biodex Medical Systems, Inc, Shirley, NY).

**Cincinnati Sportsmedicine and Orthopaedic Center Rehab Protocol: ACL – Accelerated**  
**Phase I. Weeks 1-2 (Visits: 2-4)**

<b>General Observation</b>	<ul style="list-style-type: none"> <li>v 50% weight bearing with 2 crutches when:               <ul style="list-style-type: none"> <li>- Postoperative pain controlled</li> <li>- Hemarthrosis controlled</li> <li>- Voluntary quadriceps contraction achieved</li> </ul> </li> </ul>	
<b>Evaluation</b>	<ul style="list-style-type: none"> <li>v Pain</li> <li>v Hemarthrosis</li> <li>v Patellar mobility</li> <li>v ROM minimum</li> <li>v Quadriceps contraction &amp; patella migration</li> <li>v Soft tissue contracture</li> <li>v Joint arthrometer (day 14)</li> </ul>	<p align="center"><b>Goals</b></p> Controlled Mild Good 0-110° Good None < 3 mm
<p><b>Frequency</b></p> <ul style="list-style-type: none"> <li>3-4 x/day 10 minutes</li>   <li>3 x/day 15 minutes</li>   <li>3 x/day 5 minutes</li>   <li>1-2 x/day 5 minutes</li>   <li>As required</li> </ul>	<p><b>Range of motion</b></p> ROM (passive, 0-90°) Patella mobilization Ankle pumps (plantar flexion with resistance band) Hamstring, gastroc-soleus stretches	<p align="center"><b>Duration</b></p>  5 reps x 30 secs  3 sets x 10 reps 1 set x 10 reps 3 sets x 10 reps 3 sets x 10 reps 3 sets x 10 reps 3 sets x 10 reps 3 sets x 20 reps  5 sets x 10 reps  20 minutes 20 minutes
<b>Goals</b>	<ul style="list-style-type: none"> <li>v ROM 0-110°</li> <li>v Adequate quadriceps contraction</li> <li>v Control inflammation, effusion</li> <li>v 50% weight bearing</li> </ul>	

**Cincinnati Sportsmedicine and Orthopaedic Center Rehab Protocol: ACL – Accelerated**  
**Phase 2. Weeks 3-4 (Visits: 2-4)**

<b>General Observation</b>	<ul style="list-style-type: none"> <li>v Full weight bearing with 1 crutch when:               <ul style="list-style-type: none"> <li>- Pain controlled without narcotics</li> <li>- Effusion controlled</li> <li>- ROM 0-100°</li> <li>- Muscle control throughout ROM</li> </ul> </li> </ul>	
<b>Evaluation</b>	<ul style="list-style-type: none"> <li>v Pain</li> <li>v Effusion</li> <li>v Patellar mobility</li> <li>v ROM minimum</li> <li>v Muscle control</li> <li>v Joint arthrometer (day 28)</li> <li>v Inflammatory response</li> </ul>	<p align="center"><b>Goals</b></p> <p>Mild          Minimal          Good          0-120°          3/5          &lt; 3 mm          None</p>
<b>Frequency</b>		<b>Duration</b>
3-4 x/day 10 minutes	<p><b>Range of motion</b></p> <p>ROM (passive, 0-120°)          Patella mobilization          Ankle pumps (plantar flexion with resistance band)          Hamstring, gastroc-soleus stretches</p>	5 reps x 30 secs
2-3 x/day 20 minutes	<p><b>Strengthening</b></p> <p>Straight leg raises (flexion, extension, abduction, adduction)          Isometric training: multi-angle (90, 60, 30°)          Heel raise/toe raise          Hamstring curls (active, 0-90°)          Knee extension (active, 90-30°)          Closed-chain           <ul style="list-style-type: none"> <li>- Wall sits</li> <li>- Mini-squats</li> </ul>         Multi-hip machine (flexion, extension, abduction, adduction)          Leg press (70-10°)</p>	<p>3 sets x 10 reps          1 set x 10 reps          3 sets x 10 reps          3 sets x 10 reps          3 sets x 10 reps</p>
3 x/day 5 minutes	<p><b>Balance training</b></p> <p>Weight shift side/side and forward/back          Balance board/2 legged</p>	5 sets x 10 reps
1-2 x/day 5 minutes	<p>Cup walking          Single leg stance (level surface)</p>	5 reps
2 x/day 5 minutes	<p><b>Aerobic conditioning</b></p> <p>UBC          Water walking          Stationary bicycling (patellofemoral precautions)</p>	
As required	<p><b>Modalities</b></p> <p>Electrical muscle stimulation          Cryotherapy</p>	<p>20 minutes          20 minutes</p>
<b>Goals</b>	<ul style="list-style-type: none"> <li>v ROM 0-125°</li> <li>v Muscle control</li> <li>v Arthrometer within 3 mm</li> </ul>	<ul style="list-style-type: none"> <li>v Control inflammation, effusion</li> <li>v 100% weight bearing</li> </ul>

**Cincinnati Sportsmedicine and Orthopaedic Center Rehab Protocol: ACL – Accelerated**  
*Phase 3. Weeks 5-6 (Visits: 1-2)*

<b>General Observation</b>	v Independent ambulation when: - Pain controlled                      - Effusion controlled - ROM 0-120°                              - Muscle control throughout ROM	
----------------------------	-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------	--

<b>Evaluation</b>	<ul style="list-style-type: none"> <li>v Pain</li> <li>v Effusion</li> <li>v Patellar mobility</li> <li>v ROM</li> <li>v Muscle control</li> <li>v Inflammatory response</li> <li>v Gait</li> </ul>	<b>Goals</b> No RSD Minimal Good 0-135° 4/5 None Symmetrical
<b>Frequency</b> 3 x/day 10 minutes  2 x/day 20 minutes  3 x/day 5 minutes  2 x/day 10 minutes  As required	<b>Range of motion</b> ROM (passive, 0-135°) Patella mobilization Hamstring, gastroc-soleus stretches  <b>Strengthening</b> Straight leg raises (ankle weight, not to exceed 10% of body weight) Straight leg raises, rubber tubing Isometric training: multi-angle (90, 60, 30°) Heel raise/toe raise Hamstring curls (active, 0-90°) Knee extension with resistance (90-30°) Closed-chain <ul style="list-style-type: none"> <li>- Wall sits</li> <li>- Mini-squats</li> </ul> Multi-hip machine (flexion, extension, abduction, adduction) Leg press (70-10°)  <b>Balance training</b> Balance board/2 legged Lateral step-ups: 2-4”  <b>Aerobic conditioning</b> (patellofemoral precautions) UBC Stationary bicycling Water walking Stair machine (low resistance, low stroke) Ski machine (short stride, level, low resistance) Elliptical (low resistance)  <b>Modalities</b> Cryotherapy	<b>Duration</b>  5 reps x 30 secs  3 sets x 10 reps 3 sets x 10 reps 2 sets x 10 reps 3 sets x 20 reps 3 sets x 10 reps 3 sets x 10 reps  5 reps 3 sets x 20 reps 3 sets x 10 reps 3 sets x 10 reps          20 minutes
<b>Goals</b>	<ul style="list-style-type: none"> <li>v ROM 0-135°</li> <li>v Full weight bearing, normal gait</li> <li>v Control inflammation, effusion</li> <li>v Muscle endurance</li> <li>v Recognition complications (motion loss, RSD, increased AP displacement)</li> <li>v Recognition patellofemoral changes</li> </ul>	

**Cincinnati Sportsmedicine and Orthopaedic Center Rehab Protocol: ACL – Accelerated**  
**Phase 4. Weeks 7-8 (Visits: 1-2)**

<b>General Observation</b>	<ul style="list-style-type: none"> <li>v No effusion, painless ROM, joint stability</li> <li>v Performs activities of daily living, can walk 20 minutes without pain</li> </ul>	<ul style="list-style-type: none"> <li>v ROM 0-135°</li> <li>v Full weight bearing</li> </ul>
<b>Evaluation</b>	<ul style="list-style-type: none"> <li>v Manual muscle test Hamstrings, quadriceps, hip abductors/adductors/flexors/extensors</li> <li>v Swelling</li> <li>v Joint arthrometer (8 weeks)</li> <li>v Patellar mobility</li> <li>v Crepitus</li> </ul>	<p align="center"><b>Goals</b></p> <p>4/5</p> <p>None &lt; 3 mm Good None/slight</p>
<p><b>Frequency</b></p> <p>2 x/day 10 minutes</p> <p>2 x/day 20 minutes</p> <p>3 x/day 5 minutes</p> <p>1-2 x/day 15-20 minutes</p> <p>As required</p>	<p><b>Range of motion</b> Hamstring, gastroc-soleus stretches</p> <p><b>Strengthening</b> Straight leg raises, rubber tubing Hamstring curls (active, 0-90°) Knee extension with resistance (90-30°) Leg press (70-10°) Multi-hip machine (flexion, extension, abduction, adduction) Closed-chain - Wall sits - Mini-squats</p> <p><b>Balance training</b> Balance board/2 legged Single leg stance Resistance band walking Plyoback ball toss Perturbation training</p> <p><b>Aerobic conditioning</b> (patellofemoral precautions) Stationary bicycling Water walking Swimming (straight leg kicking) Walking Stair machine (low resistance, low stroke) Ski machine (short stride, level, low resistance) Elliptical machine (low resistance)</p> <p><b>Modalities</b> Cryotherapy</p>	<p align="center"><b>Duration</b></p> <p>5 reps x 30 secs</p> <p>3 sets x 30 reps 3 sets x 10 reps 3 sets x 10 reps 3 sets x 10 reps 3 sets x 10 reps</p> <p>5 reps 3 sets x 20 reps</p> <p>20 minutes</p>
<b>Goals</b>	v Increase strength and endurance	



**Cincinnati Sportsmedicine and Orthopaedic Center Rehab Protocol: ACL – Accelerated**  
**Phase 6. Weeks 13-26 (Visits: 2-3)**

<b>General Observation</b>	<ul style="list-style-type: none"> <li>v No effusion, painless ROM, joint stability</li> <li>v Performs activities of daily living, can walk 20 minutes without pain</li> </ul>	
<b>Evaluation</b>	<ul style="list-style-type: none"> <li>v Isokinetic test (isometric + torque 300°/sec, % difference quads &amp; hams, tested monthly)</li> <li>v Swelling</li> <li>v Joint arthrometer</li> <li>v Patellar mobility</li> <li>v Crepitus</li> <li>v Single-leg function tests (hop distance, timed hop, % inv/uninv)</li> </ul>	<p align="center"><b>Goals</b></p> <p>20-25</p> <p>None</p> <p>3 mm</p> <p>Good</p> <p>None/slight</p> <p>75</p>
<b>Frequency</b>		<b>Duration</b>
2 x/day 10 minutes	<b>Range of motion</b> Hamstring, gastroc-soleus, quad, ITB stretches	5 reps x 30 secs
1 x/day 20-30 minutes	<b>Strengthening (patellofemoral precautions)</b> Straight leg raises, rubber tubing (high speed) Hamstring curls (active, 0-90°) Knee extension with resistance (90-30°) Leg press (70-10°) Multi-hip machine (flexion, extension, abduction, adduction)	3 sets x 30 reps 3 sets x 10 reps 3 sets x 10 reps 3 sets x 10 reps 3 sets x 10 reps
3 x/day 5 minutes	<b>Balance training</b> Balance board/2 legged → Single leg Single leg stance → Unstable platform	
3 x/week 20-30 minutes	<b>Aerobic conditioning</b> (patellofemoral precautions) Stationary bicycling Water walking Swimming (kicking) Walking Stair machine (low resistance, low stroke) Ski machine (short stride, level, low resistance) Elliptical machine (low resistance)	
3 x/week 15-20 minutes	<b>Running program</b> (straight) Jog (progression of speed: ½, ¾, full) Walk Backward run	1/4 – 1 mile 1/8 mile 20 yards
3 x/week 3 x/week	<b>Cutting program</b> – lateral, carioca, figure 8's <b>Functional training</b> Plyometric training: box hops, level, double-leg Sport specific drills	20 yards 15 secs, 4-6 sets
As required	<b>Modalities</b> Cryotherapy	20 minutes
<b>Goals</b>	v Increase strength and endurance	

**Cincinnati Sportsmedicine and Orthopaedic Center Rehab Protocol: ACL – Accelerated**  
**Phase 7. Weeks 27-52 (Visits: 2-3)**

<b>General Observation</b>	<ul style="list-style-type: none"> <li>v No effusion, painless ROM, joint stability</li> <li>v Performs ADL, can walk 20 minutes without pain</li> </ul>	
<b>Evaluation</b>	<ul style="list-style-type: none"> <li>v Isokinetic test (torque at 180°/sec &amp; 300°/sec, % difference quads &amp; hams, tested quarterly; isometric if symptomatic PF joint)</li> <li>v Swelling</li> <li>v Joint arthrometer</li> <li>v Patellar mobility</li> <li>v Crepitus</li> <li>v Single-leg function test (hop distance, timed hop, % inv/uninv)</li> </ul>	<p align="center"><b>Goals</b></p> <p>10-15</p> <p>None</p> <p>&lt; 3 mm</p> <p>Good</p> <p>None/slight</p> <p>85</p>
<p><b>Frequency</b></p> <p>2 x/day 10 minutes</p> <p>3-4 x/week 20-30 minutes</p> <p>3 x/day 5 minutes</p> <p>3 x/week 20-30 minutes</p> <p>3 x/week 20 minutes</p> <p>3 x/week</p> <p>3 x/week</p> <p>As required</p>	<p><b>Range of motion</b> Hamstring, gastroc-soleus, quad, ITB stretches</p> <p><b>Strengthening (patellofemoral precautions)</b>            Straight leg raises, rubber tubing (high speed)            Hamstring curls (active, 0-90°)            Knee extension with resistance (90-30°)            Leg press (70-10°)            Multi-hip machine (flexion, extension, abduction, adduction)</p> <p><b>Balance training</b>            Balance board/2 legged → Single leg            Single leg stance - unstable platform w/secondary activity</p> <p><b>Aerobic conditioning</b> (patellofemoral precautions)            Stationary bicycling            Water walking            Swimming (kicking)            Walking            Stair machine (low resistance, low stroke)            Ski machine (short stride, level, low resistance)            Elliptical (low resistance)</p> <p><b>Running program</b> (straight)            Interval training (20, 40, 60, 100 yards)            Walk/rest phase (3:1 rest: work)            Backward run</p> <p><b>Cutting program</b> – lateral, carioca, figure 8's</p> <p><b>Functional training</b>            Plyometric training: box hops, level, double-leg            Sport specific drills</p> <p><b>Modalities</b>            Cryotherapy</p>	<p align="center"><b>Duration</b></p> <p>5 reps x 30 secs</p> <p>3 sets x 30 reps</p> <p>} 1-2 sets x 8-12 reps</p> <p>20 yards</p> <p>20 yards 15 secs, 4-6 sets</p> <p>20 minutes</p>
<b>Goals</b>	<ul style="list-style-type: none"> <li>v Increase function</li> <li>v Maintain strength, endurance</li> <li>v Return to previous activity level</li> </ul>	